

Recommended Vaccines for Children from 7 to 18 Years

Child's Name:
Date of Birth:

"Recommended Vaccines for Children from 7 to 18 Years" was created and translated October 20, 2023 by the Multilingual Health Education Alliance (M-HEAL). [English]

This document is based on recommendations of the U.S. Centers for Disease Control and Prevention as of Feb 10, 2023.



If your child will be traveling outside the U.S., lives in a region where dengue infections are common, or has a health condition that can weaken the immune system, ask their doctor whether any additional vaccines or additional doses are needed to ensure protection.

If your child has not yet received all of their early childhood vaccines, see "Recommended Vaccines for Children Birth to 6 Years." Your doctor will make a different plan to ensure they get caught up on all their vaccines.

	7 Years	8 Years	9 Years	10 Years	11 Years	12 Years	13 Years	14 Years	15 Years	16 Years	17 Years	18 Years
COVID-19*	COVID-19*											
Influenza (flu)	Flu (1 or 2 doses yearly)	Flu (1 dose yearly)										
Tetanus, Diptheria, and Pertussis					Tdap							
Human papilloma virus (HPV)**			HPV (2 or 3 doses)**									
Meningococcal disease (MenACWY)					MenACWY					MenACWY		
Meningococcal disease (MenB)***										MenB***		
Additional information:	*COVID-19: Number of doses depends on age and type of vaccine used				**HPV: For children ages 9 through 12 years, there are 2 doses of HPV. For children who receive the first dose at age 16 or older, there are 3 doses of HPV.				MenB is recommended for all children at increased risk. Other children may get the vaccine if they wish.			